

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES, RUBBER BOOTS AND FACE PROTECTION (FACESHIELD) when handling contaminated surfaces and applying by hand-held equipment.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), when applying by vehicle mounted equipment.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before meals and after work.

WHEN USING DO NOT EAT DRINK OR SMOKE.

IF YOU FEEL UNWELL, seek medical advice (show the label where possible)

Storage and disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank, and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose

Environmental protection

KEEP LIVESTOCK out of treated areas for at least 14 days following treatment and until poisonous weeds such as ragwort have died and become unpalatable.

Do not contaminate surface waters or ditches with chemical or used container.

IMPORTANT INFORMATION

FOR USE ONLY AS AN HORTICULTURAL HERBICIDE

Situation	Maximum Individual Dose	Maximum Number of Treatments
Managed Amenity Turf	2.0 Litres product/ha	Up to 2 per year

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.



Holster XL[®]

MAPP No. 13596

Holster XL is an emulsifiable concentrate containing 105 g/l (9.76 % w/w) fluroxypyr as the 1-methylheptyl ester, 285 g/l (26.5% w/w) 2,4-D as the 2-ethylhexyl ester and 52.5 g/l (4.88% w/w) dicamba.

A selective post-emergence herbicide for the control of a range of broad-leaved weeds in Managed amenity turf including fine turf.

**HARMFUL IF SWALLOWED
IRRITATING TO EYES AND SKIN
MAY CAUSE LUNG DAMAGE IF SWALLOWED**

To avoid risks to man and the environment, comply with the instructions for use.

Safety data sheet available for professional user on request.

This product is approved under The Control Of Pesticide Regulations 1986 (as amended).

The Control of Substances Hazardous to Health (COSHH) Regulations may apply to the use of this product at work.

PROTECT FROM FROST



HARMFUL

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

MODE OF ACTION

Holster XL is a selective post-emergence herbicide for the control of a range of broad-leaved weeds in Managed amenity turf including fine turf.

USES

All varieties of turf may be treated, newly sown turf must only be treated in the Spring providing that at least two months have elapsed between sowing and application.

Holster XL has been tested for selectivity on the following turf grass species:

Perennial Rye-Grass	Chewings Fescue
Annual Meadow-Grass	Slender Creeping Red Fescue
Bents	Smooth-Stalked Meadow-Grass

WEEDS CONTROLLED

Perennial Weeds

Cat's-ear	Mouse-ear Hawkweed
Creeping Buttercup	Plantain spp
Common mouse-ear	Selfheal
Common Sorrell	Slender Speedwell
Daisy	White Clover
Dandelion	Yarrow

Annual Weeds

Common Chickweed	Lesser Trefoil (Yellow Suckling Clover)
------------------	---

Marketing Company: Barclay Chemicals Manufacturing Ltd., Damastown Way, Damastown Industrial Park, Mulhuddart, Dublin 15, Ireland.

Tel: 00 353 (0) 1 8112900 Fax: 00 353 (0) 1 8224678 E-mail: info@barclay.ie Website: www.barclay.ie

Approval Holder: Barclay Chemicals R&D Ltd

Contact details as above.

Copyright © Barclay Chemicals (R&D) Limited, 2007.

Holster XL is a registered trademark of Barclay Chemicals (R&D) Ltd



Holster XL

APPLICATION

Apply as a MEDIUM spray taking care to avoid spray drift.

Water Volume

Apply the recommended rate of Holster XL in the amount of water indicated in the table. Use clean water. Dirty or muddy water may reduce effectiveness

Vehicle mounted sprayer-	Holster XL should be applied at 2.0 l/ha-	200-1000 l/ha
Knapsack and hand held sprayers-	Holster XL should be applied at 20ml in 10 litres of water per 100m ²	

Number of applications

Two applications may be made per year

Timing

Apply when weeds are actively growing, normally between April and September and when the soil is moist. Do not spray if turf is wet. Do not spray in drought conditions or if turf is under stress from frost, waterlogging, trace element deficiency, pest or disease attack. Do not spray if frost is imminent or when periods of prolonged cold weather or low overnight temperatures occur. Avoid mowing for 3 days before and after spraying. Repeat application 8-10 weeks later if necessary.

Best results will be obtained by application in spring or early summer before weeds begin to flower. Avoid drift onto non-target crops, as very small quantities may cause severe damage. Avoid overlap or accidental overdosing especially on newly sown turf.

Mixing

Shake container well, half fill the spray tank with clean water. Add the required quantity of Holster XL to the spray tank and mix thoroughly. Top up the spray tank with water to the required level. Agitate thoroughly before use.

Care of sprayer

After each days use, wash out the sprayer thoroughly with water and a wetting agent at the recommended rate of use.