




Holster XL™

Holster XL is a selective post-emergence herbicide for the control of a range of broad-leaved weeds in managed amenity turf including fine turf and amenity grassland

Contains 105 g/l (9.76 % w/w) fluroxypyr as the 1-methylheptyl ester, 285 g/l (26.5% w/w) 2,4-D as the 2-ethylhexyl ester and 52.5 g/l (4.88% w/w) dicamba as an Emulsifiable Concentrate.

Safety Information **Danger**

Harmful if swallowed
May be fatal if swallowed and enters airways
Causes skin irritation
Causes serious eye irritation
May cause drowsiness or dizziness
Very toxic to aquatic life
Very toxic to aquatic life with long lasting effects
If SWALLOWED: Rinse mouth. Do NOT induce vomiting
If eye irritation persists: Get medical advice/attention.
Call a POISON CENTRE / doctor if you feel unwell.
Store in a well-ventilated place. Keep container tightly closed.
Avoid release to the environment.
Dispose of contents/container to a licensed waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.
To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies
To avoid risks to human health and the environment, comply with the Instructions for use.



PCS No. 04854

Holster XL is a selective post-emergence herbicide for the control of a range of broad-leaved weeds in managed amenity turf including fine turf and amenity grassland

Manufacturer:

Barclay Chemicals Manufacturing Ltd.,
Damastown Way,
Damastown Industrial Park,
Mulhuddart, Dublin 15, Ireland
Tel: +353 1 8112900
Fax: +353 1 8224678
E-mail: info@barclay.ie
Website: www.barclay.ie

Approval Holder:

Barclay Chemicals (R&D) Ltd.
Contact details as above.

Copyright © Barclay Chemicals (R&D) Ltd.
2016.

TMHolster is a trademark of
Barclay Chemicals (R&D) Ltd

PROTECT FROM FROST

*Contains 105 g/l
(9.76 % w/w) fluroxypyr as
the 1-methylheptyl ester,
285 g/l (26.5% w/w)
2,4-D as the 2-ethylhexyl
ester and 52.5 g/l
(4.88% w/w) dicamba as
an Emulsifiable
Concentrate.*

**FOR USE ONLY AS
AN AGRICULTURAL/
HORTICULTURAL
HERBICIDE
(Please see inside
for DIRECTIONS
FOR USE)**

Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

In case of emergency contact the Poisons Information Centre
Tel: +353 1 8092566 or +353 1 8379964

DIRECTIONS FOR USE

The following table pertains to and forms part of the STATUTORY CONDITIONS RELATING TO USE

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE				
Crop or situation	Maximum individual dose of product	Maximum number of treatments	Maximum total dose	Latest time of application
Managed amenity turf, amenity grassland	2.0 litres product per hectare	2 per year	4.0 litres product per hectare per year	N/A
New-sown amenity grassland, new-sown managed amenity turf	0.95 l/ha	Up to 2 per year	1.9 l/ha product per year	N/A

GENERAL INFORMATION

Holster XL is a selective post-emergence herbicide for the control of a range of broad-leaved weeds in managed amenity turf, including fine turf, and amenity grassland. To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

TIMING

Best results will be obtained by application in spring or early summer before weeds begin to flower. Avoid spray drift onto neighbouring crops and other plants outside the target area. Do not spray in windy weather. Beets, all brassicae including oilseed rape, lettuce, peas, potatoes, tomatoes, all fruit crops and ornamentals are particularly susceptible to Holster XL and may be severely damaged by any spray drift. Avoid overlap or accidental overdosing especially on newly sown turf.

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered to be resistant to a herbicide if it survives a correctly-applied treatment at the recommended dose. A strategy for preventing and managing such resistance should be adopted. This should include integrating herbicides with a programme of cultural control measures. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer.

WEATHER CONDITIONS

Do not spray if turf is wet. Do not spray in drought conditions or if turf is under stress from frost, water logging, trace element deficiency, pest or disease attack. Do not spray if frost is imminent or when periods of prolonged cold weather or low overnight temperatures occur. Repeat application 8-10 weeks later if necessary (minimum interval 21 days between applications).

FOLLOWING CROPS

Do not sow any crop into soil treated with HOLSTER XL for at least 3 months after application. All crop residues derived from grassland treated with HOLSTER XL at more than 2 l/ha in total in any one year must be returned to grassland or land to be cropped with cereals or forage maize.

MIXING

Shake container well, half fill the spray tank with clean water. Add the required quantity of Holster XL to the spray tank and mix thoroughly. Top up the spray tank with water to the required level. Agitate thoroughly before use.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

APPLICATION & SPRAY QUALITY

Apply the recommended rate of Holster XL in the amount of water indicated in the table. Use clean water. Dirty or muddy water may reduce effectiveness.

CARE OF EQUIPMENT

Directly after each days use with HOLSTER XL, wash out the sprayer thoroughly with clean water and a wetting agent recommended for the cleaning of sprayers. Any traces of herbicide left in the sprayer may damage susceptible crops when the equipment is subsequently used.

MODE OF USE:

Holster XL is a selective post-emergence herbicide for the control of a range of broad-leaved weeds in managed amenity turf, including fine turf, and amenity grassland.

CONTROL OF BROADLEAVED ANNUAL AND PERENNIAL WEEDS

All varieties of turf and amenity grassland may be treated; newly sown turf must only be treated in the Spring providing that at least two months have elapsed between sowing and application.

Holster XL has been tested for selectivity on the following turf grass species:

Perennial Rye-Grass	Chewings Fescue
Annual Meadow-Grass	Slender Creeping Red Fescue
Bents	Smooth-Stalked Meadow-Grass

Weeds Controlled at 2 L/ha:

Established Perennial Weeds in established managed amenity turf or amenity grassland

Cat's-ear	Mouse-ear Hawkweed
Creeping Buttercup	Plantain spp
Common mouse-ear	Selfheal
Common Sorrell	Slender Speedwell
Daisy	White Clover
Dandelion	Yarrow
Docks	

Annual Weeds in established managed amenity turf or amenity grassland

Common Chickweed	Lesser Trefoil (Yellow Suckling Clover)
------------------	---

Weeds Controlled at 0.95 l/ha

Annual Weeds in established managed amenity turf or amenity grassland

Common Chickweed MS

APPLICATION
ESTABLISHED MANAGED AMENITY TURF OR AMENITY GRASSLAND

Crop	Time and method	Dose rate
Managed amenity turf Amenity grassland	Apply when weeds are actively growing, normally between April and September and when the soil is moist. Using a vehicle mounted sprayer	2.0 l/ha Apply in 200-1000 l/ha water, as a MEDIUM quality spray
Managed amenity turf	Apply when weeds are actively growing, normally between April and September and when the soil is moist. Using a knapsack sprayer	20ml in 10 litres of water, to treat 100m ²

NEW-SOWN MANAGED AMENITY TURF OR AMENITY GRASSLAND

Crop	Time and method	Dose rate
Managed amenity turf Amenity grassland	Apply when weeds are actively growing, in Spring at least 2 months after grass emergence and when the soil is moist. Using a vehicle mounted sprayer	0.95 l/ha Apply in 200-1000 l/ha water, as a MEDIUM quality spray
Managed amenity turf	Apply when weeds are actively growing, in Spring at least 2 months after grass emergence and when the soil is moist. Using a knapsack sprayer	9.5ml in 10 litres of water, to treat 100m ²